

NCI-H1299 | 300485

<b>Description</b>	H1299	NCI-H1299	43		. H1299	H292	(NSCLC)	.
		, H1299	p53		p53		. KRAS	
		KRAS			.			.
	H1299	5		. H1299	22~30	2	. H1299	
		B(NMB) 0.1 pmol/mg			(GRP)		.	A549

H1299

**Organism**

**Tissue**

**Disease**

**Synonyms**

H1299, H-1299, NCH1299

**Age**

59

**Ethnicity**

**Growth properties**

**Citation**

NCI-H1299( 300485)

**Biosafety level**

1

**NCBI\_TaxID**

9606

**CellSaurusAccession**

CVCL\_0060

NCI-H1299 | 300485

**Culture Medium** RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3( 820700a)

**Supplements** 10% FBS 2.5g/L 10mM HEPES

**Dissociation Reagent**

**Subculturing** PBS . T25 3~5ml, T75 5~10ml PBS . T25

**Fluid renewal** 2~3

**Freeze medium** (FBS ) + 10% DMSO , CM-1(Cytion 800100)

**Thawing and Culturing Cells**

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

**Incubation Atmosphere** 37°C, 5% CO2' .

**Flask Coating**

NCI-H1299 | 300485

**Freezing  
Procedure**

-78°C

**Shipping  
Conditions**

-78°C

**Storage  
Conditions**

-150°C -196°C .80°C

/ / HLA

**Sterility**

PCR