

HEK293-FAP | 305419

HEK293-FAP

Culture Medium RPMI 1640, w: 2.0 mM β -mercaptoethanol, w: 2.0 mM/1000 NaHCO₃ (Cytion 820700a)

Supplements FBS 10%, β -mercaptoethanol 1 mM, HEPES 10 mM, NEAA 1% (Cytion 820700a)

Dissociation Reagent Trypsin-EDTA

Subculturing 1:2 to 1:10 in fresh medium

Fluid renewal 2-3 times per week

Post-Thaw Recovery Seed cells into T25 flasks at 1:2 to 1:3 dilution in fresh medium

Freeze medium DMEM (Cytion 820700a) + 10% FBS + 10% DMSO

- Thawing and Culturing Cells**
1. Thaw cryovial in 37°C water bath
 2. Dilute cells into 10 ml DMEM + 10% FBS
 3. Seed cells into T25 flask
 4. Incubate cells at 37°C in 5% CO₂ for 24 hours
 5. Refresh medium after 24 hours
 6. Harvest cells after 3-5 days
 7. Seed cells into T25 flask at 1:2 to 1:10 dilution
 8. Incubate cells at 37°C in 5% CO₂

Product sheet

HEK293-FAP | 305419

Incubation Atmosphere 37°C, 5% CO₂, [REDACTED]

Flask Coating [REDACTED]

Freezing Procedure [REDACTED]

Shipping Conditions [REDACTED]

Storage Conditions [REDACTED] -150 [REDACTED] -196 °C [REDACTED]

[REDACTED] / [REDACTED] / HLA

Sterility [REDACTED] PCR [REDACTED]
[REDACTED] [REDACTED] [REDACTED]